

LA-UR -80-46

TITLE: TWO-SECTION TRANSFER STUDY OF ^{195}Pt

AUTHOR(S): J. A. Cizewski, P. R. Flynn, Donald L. Brown, and L. W. Simler

SUBMITTED TO: Conference on Band Structure and Nuclear Dynamics,
New Orleans, LA, February 28 - March 1, 1980

MASTER

University of California



LOS ALAMOS SCIENTIFIC LABORATORY

Post Office Box 1663 Los Alamos, New Mexico 87545

An Affirmative Action/Equal Opportunity Employer

By acceptance of this article, the publisher recognizes that the
U.S. Government retains a nonexclusive right to reproduce
or publish in reproduction the published form of this content
herein, or to allow others to do so, for U.S. Government purposes.

The Los Alamos Scientific Laboratory reports that the publisher
hereby identifies this article as work performed under the auspices
of the U.S. Department of Energy.

TWO-NEUTRON TRANSFER STUDY ON ^{195}Pt .*

J. A. Sizewski, E. R. Flynn, Ronald E. Brown, and J. W. Sunier
Los Alamos Scientific Laboratory, Los Alamos, NM 87545

ABSTRACT

We have investigated the $^{195}\text{Pt}(t,p)^{197}\text{Pt}$ reaction. Three states below 1 MeV are populated with a large fraction of the $L=0$ strength, which is seen in the even to even cases. This is in contrast to other (p,t) and (t,p) studies on Pt targets, where the $L=0$ strength is concentrated in the ground state.

INTRODUCTION

In two-particle transfer studies one expects to populate strongly states with wave functions that have a large overlap with the target ground state. In odd-A nuclei, where the odd particle is weakly coupled to the even-even core, the (p,t) or (t,p) reaction can be used to identify the core-coupled state,¹ since the population pattern will be similar to that of the (p,t) or (t,p) reaction on the even-even core after blocking corrections. Alternatively, two-particle transfer studies are a sensitive probe of transitional regions, because of the fragmentation of $L=0$ transfer strength observed in these regions. For example, in a spherical-deformed transitional region² the levels populated in the final deformed nucleus will be states described by Nilsson orbitals having large components of the spherical-target, ground-state orbital. To probe the A 190 transitional region, we have initiated an investigation of the (t,p) reaction systematically starting with the ^{195}Pt target.

EXPERIMENTAL PROCEDURES

The $^{195}\text{Pt}(t,p)$ reaction was investigated with a 17-MeV triton beam from the IAS1 tandem Van de Graaff accelerator. The reaction protons were detected by a helical proportional counter³ in the focal plane of the Q3D spectrometer.⁴ The levels in ^{197}Pt below 1 MeV populated in the present study are indicated by an X in Figure 1, which summarizes the information known⁵⁻⁸ about low-lying negative-parity levels in $^{193,195,197}\text{Pt}$. The $1/2^-$ states in ^{197}Pt at 0, 131 and 748 keV are populated by $L=0$ transitions with cross sections at 25° of 103, 20, and 30 $\mu\text{b/sr}$, respectively. In Table 1 we summarize the systematics^{8,9} of relative ground-state, $L=0$ transitions from $\text{Pt}(t,p)$ and (p,t) reaction studies. We have also indicated the summed $L=0$ strength below 1 MeV for $^{195}\text{Pt}(t,p)$.

DISCUSSION

Figure 1 indicates the differences between ^{197}Pt and $^{193,195}\text{Pt}$. In ^{197}Pt three low-lying $1/2^-$ states have been identified, as opposed to only one in $^{195,193}\text{Pt}$, and all three of these

TABLE I: Relative Ground-State Transition Strengths for Two Neutron Transfer Studies on Pt Nuclei.^{a)}

	TARGET				
	¹⁹² Pt	¹⁹⁴ Pt	¹⁹⁵ Pt	¹⁹⁶ Pt	¹⁹⁸ Pt
(p,t)	111	100	50	97	98.5
(t,p)	92	100	28 (42)	97	88

a) The data are from ref. 8, 9 and the present study. For the ¹⁹⁵Pt target the summed l.u. strength below 1 MeV is given in parentheses.




Figure 1. Negative-parity states in ¹⁹²Pt, ¹⁹⁴Pt, ¹⁹⁶Pt.

states have wave functions with sizeable overlap with the ^{195}Pt ground state, since they are all populated quite strongly in the (t,p) reaction. In the even Pt(t,p) and (p,t) reactions⁸⁻¹⁰ almost all of the $L=0$ strength goes to the ground state (as is also the case for $^{195}\text{Pt}(p,t)$), with no excited state receiving more than 10% of the ground-state strength, and excited 0^+ states typically are above 1 MeV in excitation. Therefore, the $1/2^-$ states in ^{197}Pt are not simple core-coupled excitations.

The Pt nuclei are in an intermediate region between well-deformed and spherical nuclei. ^{195}Pt has frequently⁶ been described within an oblate framework, but if ^{197}Pt were less deformed than ^{195}Pt , one would expect fewer rather than more $1/2^-$ states. In fact, in a Nilsson-model description one cannot account for three low-lying $1/2^-$ states. The fragmentation of single-particle strength in Hf-W-Os nuclei has recently been understood as resulting from changes in the hexadecapole and quadrupole deformations of these nuclei. The even-even Pt cores, however, are exhibiting a quite stable structure⁹, except possibly for $^{198,200}\text{Pt}$. Therefore, traditional approaches such as the Nilsson model, even accounting for fragmentation of single-particle strength as was done in ref. 11, seem to be inadequate to describe the ^{197}Pt level structure. A more complete investigation of the (t,p) and (p,t) reactions in the A 190 nuclei, especially with N 120, and additional knowledge of ^{199}Pt , will be necessary before a good understanding of these nuclei will be possible.

REFERENCES

*Work supported by The U. S. Department of Energy.

1. For example, R. E. Anderson, et al., Phys. Rev. C15, 123 (1977).
2. For example, P. G. Burke, et al., Nucl. Phys. A 258, 113 (1976).
3. L. R. Flynn, Nucl. Instr. and Meth. 162, 305 (1974), and references therein.
4. L. R. Flynn, et al., Nucl. Instr. and Meth. 129, 35 (1975).
5. G. R. Smith, et al., Nucl. Phys. A 290, 72 (1977); G. Berrier-Ronsin, et al., Phys. Rev. C17, 529 (1978); and references therein.
6. Y. Yamazaki and R. E. Sheline, Phys. Rev. C14, 531 (1976), and references therein.
7. Y. Yamazaki, et al., Phys. Rev. C17, 2061 (1978), and references therein.
8. G. Rothbard, et al., Orsay report IPNO PhN 79-1, (1979), and references therein.
9. J. A. Cizewski, et al., Phys. Lett., and to be published.
10. P. T. Deason, et al., Phys. Rev. C20, 922 (1979).
11. R. E. Casten, et al., Nucl. Phys., to be published.